

Leptodea fragilis (Raf.)	Lampsilis ovata ventricosa
Proptera alata (Say)	(Barnes)
Micromya ortmanni Walker	Truncilla truncata Raf.
Lampsilis siliquoidea (Barnes)	Dysnomia triquetra (Raf.)
1924	1925
Lampsilis ovata (Say) 1925	

BIBLIOGRAPHY

- 1895—Cooke, A. H. Camb. Nat. Hist. Mollusca.
 1920—Grier, N. M. NAUTILUS, Vol. XXXIV.
 1922—Jewell, M. E. Ecology, Vol. III.

 A NEW PECTEN FROM THE LOOCHOO ISLANDS

BY W. H. DALL

PECTEN (CHLAMYS) SMEGMATUS n. sp.

Shell compressed, glistening, whitish above with reddish flecks, darker and more crowded distally; a few opaque white zigzag markings near the umbo; lower valve white with purplish flush near the umbo and reddish margin on the wings; sculpture of the upper valve with nine low subangular whitish rays with one smaller ray in the interspaces with one or two small threads on each side of it; wings nearly smooth, oblique, the left wing much the largest; sculpture of the lower valve of numerous low undulated riblets with one or two small threads in the interspaces; the sculpture grows feeble toward the umbo; byssal notch wide and deep, ctenolium with eleven denticulations; height of shell, 33; width of shell, 30; of hinge-line, 17; diameter, 7 mm.

Loochoo Islands; Coll. Thaanum, No. 9455.

This is nearest some of the forms of *Pecten squamosus* Gmelin, but has a different sculpture with no indication of raised scales. The surface has a greasy sheen. A single specimen was obtained by Messrs. Thaanum and Langford, during their collecting trip of 1925.